Review Article



State of the structure address on MET receptor activation by HGF

[©] Edmond M. Linossi¹, Gabriella O. Estevam², Masaya Oshima^{4,5}, James S. Fraser², Eric A. Collisson^{4,5} and Natalia Jura^{1,3}

¹Cardiovascular Research Institute, University of California – San Francisco, San Francisco, CA 94158, U.S.A.; ²Department of Bioengineering and Therapeutic Sciences, University of California – San Francisco, San Francisco, CA 94158, U.S.A.; ³Department of Cellular and Molecular Pharmacology, University of California – San Francisco, San Francisco, CA 94158, U.S.A.; ⁴Division of Hematology and Oncology, Department of Medicine, University of California, San Francisco, CA 94158, U.S.A.; ⁵UCSF Helen Diller Family Comprehensive Cancer Center, University of California - San Francisco, San Francisco, CA 94158, U.S.A.

Correspondence: Edmond M. Linossi (edmond.linossi@ucsf.edu) or Natalia Jura (natalia.jura@ucsf.edu)

The MET receptor tyrosine kinase (RTK) and its cognate ligand hepatocyte growth factor (HGF) comprise a signaling axis essential for development, wound healing and tissue homeostasis. Aberrant HGF/MET signaling is a driver of many cancers and contributes to drug resistance to several approved therapeutics targeting other RTKs, making MET itself an important drug target. In RTKs, homeostatic receptor signaling is dependent on auto-inhibition in the absence of ligand binding and orchestrated set of conformational changes induced by ligand-mediated receptor dimerization that result in activation of the intracellular kinase domains. A fundamental understanding of these mechanisms in the MET receptor remains incomplete, despite decades of research. This is due in part to the complex structure of the HGF ligand, which remains unknown in its full-length form, and a lack of high-resolution structures of the complete MET extracellular portion in an apo or ligand-bound state. A current view of HGF-dependent MET activation has evolved from biochemical and structural studies of HGF and MET fragments and here we review what these findings have thus far revealed.

Introduction

MET, also known as hepatocyte growth factor receptor (HGFR) and its close homolog RON (Recepteur d'Origine Nantais), also known as macrophage-stimulating 1 receptor (MST1R) form a subfamily within the larger receptor tyrosine kinase (RTK) family of membrane receptors [1–5]. Like other RTKs, MET consists of an extracellular domain (ECD), a single-pass transmembrane domain (TM), and an intracellular region that includes a juxtamembrane domain (JM), kinase domain (KD) and carboxy-terminal tail (C-tail) [5]. The ECD of MET includes a sema domain, a plexin/sema-phorin/integrin (PSI) domain, and four integrin/plexin/transcription (IPT) factor domains [6,7]. The single-pass transmembrane helix of MET connects the ECD module to a long intracellular juxtamembrane region that is followed by a tyrosine KD and a short C-tail (Figure 1A,B). The broad range of physiological outputs downstream from MET includes activating cell motility, altering cellular morphology and driving cell cycle progression, ultimately playing essential roles in the development of multiple tissues during embryogenesis and in wound healing and tissue homeostasis in adults [8–10]. The growth and motility signals activated by MET make it a potent oncogene when it becomes abnormally activated through gene amplification, enhanced expression or mutation, and this has been observed in many solid cancers, including lung, kidney and liver [11–13].

Received: 22 December 2020 Revised: 22 February 2021 Accepted: 24 February 2021

Version of Record published: 16 April 2021

The cognate ligand for MET is hepatocyte growth factor (HGF), also known as scatter factor (SF) [14–17]. HGF is produced in a pro-form that is proteolytically cleaved to a mature, active α - and β -chain heterodimer held together by a disulfide bond [18] (Figure 1C). Two natural splice variants of HGF, called NK1 and NK2, encompass different fragments of the HGF α -chain (Figure 1C) [19,20].





Figure 1. Architecture of the MET extracellular domain.

(A) Domain architecture of the MET receptor which includes the sema domain, PSI (Plexin, Semaphorin, Integrin) domain, IPT (Integrin, Plexin, Transcription factor) domain, TM (transmembrane) domain, JM (juxtamembrane) domain and KD (kinase domain). Residues at domain boundaries are marked. The furin cleavage site in the sema domain is indicated with a scissors sign. (B) Cartoon representation of the domain structure of the full-length MET receptor colored relative to (A). The extracellular domain (ECD) includes the sema domain and the stalk region (PSI-IPT₁₋₄). Numbers in the sema domain denote individual blades of the β -propeller structure. Exon 14 in the membrane-proximal region of the JM domain is marked by a yellow box. (C) Domain architecture of MET ligands: HGF, NK1, NK2 and InIB. The HGF cleavage site, which generates the mature α- and β-chains of the ligand, is marked with a scissors sign. Domain abbreviations are: N (N domain), K1-K4 (kringle domains 1-4), SPH (serine protease homology domain), Cap (Cap domain), LRR (leucine-rich repeat), IR (inter repeat region), B (B-repeat), GW (Gly-Trp domain). Domains colored in white have no associated high-resolution structural data. Panel on the right illustrates HGF ligand cleavage and proposed associated conformational changes that lead to formation of an active ligand. The N, K1, and K2 domain arrangement is based on crystal structures described in Figure 4A,B. (D) Crystal structure of the MET sema domain is shown in cartoon representation (PDB: 1SHY). PSI and SPH domains are removed for clarity. The extrusion region is colored lime green. (D, lower inset) The 4 antiparallel β-strands (labeled a-d) that constitute each blade of the sema domain are shown for blades 1 and 7. Strand d of blade 7, which is contributed by the N-terminal region of the sema domain, encloses the structure. (D, high inset) The extrusion region is highlighted by rotation of the sema domain to demonstrate its continuation from blade 5c that continues back to form blade 4e. Two colored dots indicate an unresolved region of the extrusion (Thr 402-Arg 413). (E) Two orientations of the MET ECD in complexes with different ligands illustrate flexibility of the stalk region. Left, structure of MET in the MET/InIB complex (PDB: 6GCU) in which the InIB ligand has been removed for clarity. Right, structure of MET in the MET/HGF^{SPH} complex (PDB: 1SHY), in which the HGF^{SPH} domain has been removed for clarity, is also aligned via the PSI domain on the structure of the PSI-IPT₁₋₂ module (PDB: 5LSP) to highlight the extended conformation of the stalk in this state. (F) The structures of MET/InIB and MET/HGF^{SPH} (shown in gray) were aligned on their sema domains and rotated as noted relative to (E). HGF^{SPH} and InIB ligands were removed for clarity.



The prevailing model for MET activation is that the mature HGF ligand can dimerize and bind two MET molecules to form a 2:2 heterocomplex [21–23]. While not strictly required for signaling, heparan sulfate can augment MET activation by HGF by driving ligand dimerization or through increasing the local concentration of HGF at the cell surface [24,25]. In addition to HGF, many other non-canonical mechanisms of MET activation have been described under pathological and physiological conditions. Internalin B (InIB), an effector protein from Listeria monocytogenes, can directly engage MET receptor forming oligomers that in turn cluster MET, activating it and triggering endocytosis that facilitates entry of the bacterium into cells [26-28]. Another protein ligand described to interact with MET is decorin, a leucine-rich repeat (LRR) proteoglycan, which upon MET binding induces receptor down-regulation through enhanced recruitment of the E3-ligase Cbl [29]. In addition, numerous reports have documented co-activation of MET via other transmembrane receptors, including Integrinβ-4 [30], CD44v6 [31], VEGFR [32], ROR1 [33], HER3 [34], HER2 [35], AXL [36], RON [37] and the plexin-B1/SEMA4D complex [38] (reviewed in [39]). The mechanisms by which these co-receptors activate MET have not been structurally defined, and in many cases, it remains unclear whether they represent direct interactions and how they converge on the canonical receptor activation mechanism that proceeds via MET homo-dimerization. Signaling cross-talk of MET with other receptors often manifests under pathological conditions and this may present novel therapeutic opportunities for targeting MET [12,40,41].

Our knowledge of activation mechanisms and communication across the plasma membrane for many RTKs has been compiled from high-resolution structures of individual domains of the receptors, their biochemical characterizations and cell-based functional studies [5,42]. While several high-resolution structures of fragments of the MET ECD and HGF have been solved (Tables 1 and 2), many important structural states are still missing. Consequently, our understanding of how HGF engages with the MET ECD to initiate activation of the intracellular KD remains incomplete. Here, we discuss the current state of structural understanding of HGF-dependent MET activation and highlight areas for much needed future investigations.

Extracellular domain of MET and its relations to the plexin/ semaphorin system

One of the distinct features of MET, and its close homolog RON, compared with other RTKs is the presence of a sema domain within their ECDs. Sema domain is a characteristic feature of another class of membrane receptors, called plexins, and their ligands, semaphorins [43]. Instead of the KD, plexins have intracellular GTPase Activating Protein (GAP) domain as signaling units [44]. It is presumed that MET and plexins shared a common ancestor, as early MET-like orthologs identified in both Lophotrochozoa and Echinodermata have extended ECDs like plexins [45], but the ligand systems for MET and plexins have evolved to be quite different. The

		MET domains						
PDB ID	Technique	sema	PSI	IPT1	IPT2	IPT3	IPT4	Other ligands/proteins
1SHY	X-ray	+	+					HGF β-chain
2UZX	X-ray	+	+	+	+			InIB
2UZY	X-ray	+	+	+	+			
403T	X-ray	+	+					HGF β -chain + Zymogen Activator Peptide (ZAP.14)
403U	X-ray	+	+					HGF β -chain + Zymogen Activator Peptide (ZAP2.3)
6GCU	X-ray	+	+	+	+			InIB and DARPin A3A
6104	X-ray	+	+					Fab
4K3J	X-ray	+	+					HGF β-chain + Fab (Onartuzumab)
1SSL	NMR		+					-
5LSP	X-ray			+	+			Fab (107_A07)
-	cryo-EM / SAXS	+	+	+	+	+	+	Low resolution maps of full MET ECD in the presence and absence of cleaved and uncleaved full length HGF from single particle analysis by cryo-EM and correlation with SAX envelopes (models not available)

Table 1. Structures of the MET ECD



		HGF domains						
PDB ID	Technique	N	K1	K2	К3	K 4	SPH	Other ligands/proteins
6LZ9	X-ray						+	Fab (t8E4)
4D3C	X-ray	+	+					monoclonal antibody (SFN68)
5CP9	X-ray	+	+					MB605 (3-(furan-2-yl)propanoic acid)
5COE	X-ray	+	+					HEPES
5CS5	X-ray	+	+					PIPES
5CS9	X-ray	+	+					MES
5CS1	X-ray	+	+					-
5CS3	X-ray	+	+					(H)EPPS
5CT1	X-ray	+	+					CHES
5CT3	X-ray	+	+					2FA (3-hydroxypropane-1-sulfonic acid)
5CSQ	X-ray	+	+					MOPS
5CT2	X-ray	+	+					CAPS
403T	X-ray						+	ZAP.14
403U	X-ray						+	ZAP2.3
4K3J	X-ray						+	Fab (Onartuzumab) + MET sema + PSI
4IUA*	X-ray	+	+	+				*Mouse NK2
3SP8	X-ray	+	+	+				Heparin DP10
ЗМКР	X-ray	+	+					Heparin
3HMS	X-ray	+						-
3HN4	X-ray	+	+	+				HEPES
ЗНМТ	X-ray	+						-
2QJ4*	X-ray	+	+					*Mouse NK1
2QJ2	X-ray	+	+					-
1SI5	X-ray						+	-
1SHY	X-ray						+	MET sema + PSI
1GP9	X-ray	+	+					HEPES
1GMN	X-ray	+	+					Heparin, HEPES
1GMO	X-ray	+	+					Heparin, HEPES
1NK1	X-ray	+	+					-
1BHT	X-ray	+	+					HEPES
2HGF	NMR	+						-
-	cryo-EM / SAXS	+	+	+	+	+	+	Low resolution maps of single chain (uncleaved) and two chain (cleaved) HGF from single particle analysis by cryo-EM and correlation with SAX envelopes (models not available)

Table 2. Structures of HGF

semaphorins commonly act as sema domain-mediated dimers to activate plexins [46] (Figure 2A). While plexins have their own sema domains, they do not dimerize via these domains but use them to bind semaphorin dimers to form an active tetrameric complex [46] (Figure 2B). The sema domain of MET also provides the predominant binding site for its ligand HGF and like semaphorins, HGF also dimerizes to bring together two MET monomers. However, since HGF does not contain a sema domain the mode of HGF binding to MET is unique.

The sema domain of MET forms a canonical β -propeller comprised of seven blades, with each blade containing 4-antiparallel β -strands named a-d moving from the inside to the outside of the propeller structure (Figure 1B,D). The circular arrangement of the seven blades results in an enclosed structure, with the





Figure 2. Architecture of the semaphorin/plexin ECDs.

(A) Surface representation of the Sema6A dimer and a PlxnA2 monomer based on PDB: 30KY. The tetrameric complex of the Sema6A dimer bound to two PlxnA2 monomers is shown boxed on the right-hand side. (B) Model of the complete MET ECD based on low resolution structures [21,58]. The MET sema, PSI and IPT1–2 are shown in surface representation from PDB: 1SHY and 5LSP. The IPT3 and IPT4 domains have not been resolved in high resolution and are shown as IPT domain silhouettes to match low resolution MET ECD models. The crystal structure of the full PlxnA1 ECD (PDB: 5L56) is also shown as a surface representation for comparison to the shorter MET ECD model.

N-terminus providing the last strand of the seventh blade (Figure 1D, inset). Loops connecting a-b and c-d form the bottom face and b-d and d-a strands form the top face of the propeller (Figure 1D). These loops represent the most divergent sequences in sema domain-containing proteins and are frequently utilized for protein-protein interactions [43,46]. Of note is the extended loop that follows the 5c strand, termed the extrusion [47], which in semaphorins forms part of the dimer interface [47–51]. In MET this loop consists of 57 amino acids and extends to form a fifth strand on the fourth blade (4e) before continuing to strand 6a, resulting in only three strands in the fifth blade (5a–5c) [52] (Figure 1D). The significance of this unique extrusion in MET is not yet defined but may contribute to its distinct mode of ligand binding from the plexin receptors.

A hallmark feature of MET and RON is the cleavage of the sema domain between strands 4d and 5a (Arg 307– Ser 308 in MET) by the endoprotease furin [7,52,53]. In MET, the resulting α (~50 kDa) and β (~145 kDa) polypeptide chains remain disulfide-linked, forming the mature heterodimeric receptor (Figure 1A,B) [7]. Surprisingly, the functional relevance of the furin-mediated cleavage of MET, which is not resolved in existing crystal structures, is unknown. Mutation of this site does not appear to impede receptor maturation, ligand binding, or activation [53,54]. Given its proximity to the putative ligand-binding site for the HGF α -chain (discussed below), this site and its proteolytic processing may make currently unappreciated contribution to ligand binding.

With the exception of the viral semaphorin protein A39R, all proteins with a sema domain are followed by a PSI domain [43]. The PSI domain is a small cysteine rich knot which in MET is followed by four Ig-like IPT domains that together form the stalk region (PSI-IPT₁₋₄) [7]. In MET, two glycine residues present in the sema-PSI linker (Gly 517 and Gly 519) provide flexibility between the sema domain and the stalk, evident in different orientations that the IPT₁ domain adopts relative to the sema domain in many ligand-bound MET (and RON) ECD structures [27,52,55,56] (Figure 1E,F). The stalk region itself is more rigid in its N-terminal part due to the presence of a loop in the IPT₁ domain, termed the β -wing, that provides extended contact with the PSI domain [7,27,57] (Figure 1E). This flexible connection between the sema domain and the more rigid stalk in MET could contribute to transitions between inactive and active states of the receptor. The MET ECD in early cryo-electron microscopy (EM) tomography and small-angle X-ray scattering (SAXS) studies [21,58] (Figure 2B). The extended stalk region of plexin receptors forms an overall ring-like architecture that has been proposed to mediate receptor autoinhibition by spatially separating two receptors and preventing dimerization [59,60] (Figure 2B). Further structures of the entire MET ECD are required to define the relationship between its ECD architecture and MET autoinhibition and activation.

Ligand-induced activation

A majority of biochemical evidence and structural models derived from low-resolution techniques, such as cryo-EM and SAXS, collectively support the model in which MET activation proceeds via the formation of a heterodimeric 2:2 ligand: receptor complex [21,55,61–63]. In this model HGF dimer bridges two MET ECDs.



In contrast with receptors such as EGFR and c-KIT, whose ECDs in solution form dimeric complexes upon ligand binding [64–66], the full MET ECD bound to HGF is monomeric in solution [21]. A shorter MET ECD fragment that contains only the sema and PSI domains was shown to form a 2:2 complex with mature HGF [21] or with one of its splice variants, NK1 [62]. This suggests that the remaining region of MET ECD (corresponding to the stalk region) exerts an autoinhibitory effect on ligand-dependent ECD dimerization. In the full-length receptor, this autoinhibition may be counteracted upon ligand binding by additional dimer-stabilizing interactions contributed by the transmembrane and intracellular regions.

The architecture of the full-length, active HGF-bound MET receptor dimer is unknown. While there have been many structures of MET ECD fragments bound to β -chain of HGF, none of these structures yielded definitive dimeric complexes that explain receptor activation. The only known ligand-stabilized dimeric form of MET ECD is a complex between the sema-PSI-IPT₁₋₂ fragment of MET ECD and the InlB toxin [27,67]. InlB contains an Internalin domain, which encompasses a Cap, LRR and an internalin repeat (IR), that together comprise the minimal fragment known to efficiently activate MET in cells [68] (Figure 1C). InlB binds MET via two interfaces [27] (Figure 3A,B). As revealed by X-ray crystallography, engagement of these two sites by InlB results in a more bent MET ECD orientation compared with the one seen in the HGF β -chain-bound sema-PSI structures [27] (Figure 1E,F). In the InlB/MET ECD structures, the top face of the sema domain is predicted to tilt towards the plasma membrane, much like a wilted flower (Figure 1F).

InlB itself dimerizes via the concave face of its LRR domain and mutation of the LRR dimer interface prevents InlB-induced MET activation [27,67,69] (Figure 3C). In MET/InlB dimer, the two MET ECDs also engage via their IPT₂ domains revealing a potential dimerization mode for MET in the active dimer (Figure 3C). While the role of the IPT₂ interface for InlB or HGF-mediated MET activation has not been characterized, antibodies or engineered protein inhibitors that bind the PSI–IPT₁ and IPT₁₋₂ region of the MET stalk abrogate HGF-induced MET activation [57,70]. Thus, the PSI–IPT₁₋₂ regions are likely important for the formation of the ECD domain dimer. High-resolution structure of the full-length ECD of MET is needed to unveil the architecture of the functionally relevant active MET ECD dimer.

HGF structure and processing

HGF is a complex growth factor, which is structurally related to the serine protease plasminogen. HGF undergoes analogous processing to plasminogen from its inactive zymogen pro-form to a proteolytically cleaved (between Val 494 and Arg 495) form, which constitutes a biologically active, disulfide-linked α - and β -chain heterodimer [71,72] (Figure 1C). The α -chain is characterized by an N-terminal (N) finger or PAN domain and four kringle domains (K1–4), and the β -chain encodes a serine protease homology (SPH) domain (Figure 1C). HGF is catalytically inactive as a protease due to mutations in the canonical catalytic triad of the SPH domain [73].

The α - and β -chain of HGF contain independent binding sites for MET, and both binding events are required to activate MET. The α -chain binds MET with high affinity and can engage the receptor in both the un-cleaved and cleaved form of the ligand [71,72]. Conversely, cleavage of HGF is required for the low-affinity site located in the β -chain of HGF to bind MET and for receptor activation [71,72,74,75]. Consequently, the isolated α - and β -chains of HGF cannot independently activate MET, however, they can activate MET when combined together from isolated preparations [76]. While a crystal structure of the SPH domain bound to the MET sema domain was solved [52], there are no high-resolution structures of MET in complex with the full-length HGF α -chain alone or the NK1 and NK2 variants of HGF that correspond to two different fragments of the α -chain. Hence, major questions remain about the collective contributions of the α - and β -chains in full-length HGF to MET activation. Below we discuss key findings that emerged from the structural studies of the individual α - and β -chains of HGF about their binding to MET.

Interactions of the α - and β -chains of HGF with MET

Numerous studies support a model in which the HGF α -chain binds to β -chain fragment of the MET sema domain (blades 5–7). Deletion of the sema domain abolishes interaction of the HGF α -chain with MET and the NK1 ligand selectively cross-links with the β -chain of MET [7,61,77]. Furthermore, InIB protein and the therapeutic antibody onartuzumab, which bind to the β -chain fragment of MET sema domain, compete with HGF binding [27,78]. Lastly, mutations detected in patients that occur in the α -helix of the extrusion region (V370D and N375L) in the β -chain of MET decrease HGF affinity for MET [79,80], further substantiating that this region of the sema domain contains the binding site for HGF α -chain. Intriguingly, another primary





Figure 3. Binding modes of MET receptor ligands.

(A) Schematic representation of InIB bound to the MET ECD and (B) crystal structure of the MET ECD (shown as cartoon) bound to InIB (shown in surface mode) based on PDB: 6GCU. Two binding sites of InIB on MET are indicated by 'i' and 'ii' and are shown at higher magnification in the insets on the right. The first site (i) involves the IR domain, a low affinity binding site, which packs against two α-helices formed by the extrusion region and 5a-b loop as well as 6c-d loop of the MET sema domain. The second site (ii) is a high-affinity binding site that involves interaction between the LRR and Cap domains of InIB and the β -wing region of the MET IPT₁ domain. Numbers refer to the corresponding blades in the sema domain. (C) InIB-induced dimer of the MET ECD seen in the MET/InIB crystal structure (PDB: 6GCU) is shown (bottom) and cartooned as a schematic model (top). Dimerization interface is formed by the concave face of the LRR repeat of InIB, with a second putative interface formed by the IPT₂ domains of the MET ECD. (D) Schematic representation of HGF SPH domain bound to the MET ECD and (E) crystal structure of the MET ECD (shown as cartoon) bound to the SPH domain (shown in surface mode) based on PDB: 1SHY. The inset on the right shows a magnified view of the SPH/MET binding interface seen in the crystal structure. Upon SPH domain cleavage, the liberated SPH N-terminal peptide binds back to the activation pocket via the critical Val 495 residue engaging Asp 672 (colored in black). The pseudo-active site residues Gln 534, Asp 578, and Tyr 673 (colored yellow) together with a loop region (Val 692-Gly 696) (colored black) of the SPH domain make up the binding interface on HGF which engages the bottom face of blades 2 and 3 on the MET sema domain. (F) HGF^{SPH}-induced dimer of the MET ECD seen in the MET/HGF^{SPH} crystal structure (PDB: 1SHY) is shown (bottom) and cartooned as a schematic model (top).

binding site for HGF α -chain has been proposed by Michieli, Comoglio and colleagues in the IPT₃₋₄ domains of the MET stalk. Monoclonal antibodies targeting this region of MET do indeed reduce HGF-driven signaling [70,81]. Thus, further investigation is required to definitively resolve the α -chain binding site on MET.

The pseudo-protease (SPH domain) of the HGF β -chain is the only fragment of HGF resolved in crystal structures in complex with MET (Figure 3D,E). Like in other plasminogen proteases, the HGF SPH domain is generated via cleavage that liberates an amino-terminal polypeptide, which then binds back to the SPH domain and allosterically stabilizes the 'active site' pocket [52] (Figure 3E). The cleavage results in an approximate 14-fold increase in affinity between the HGF β -chain and MET [75]. Short peptides designed to mimic the



binding mode of the N-terminal residue of cleaved HGF to the SPH pocket increase HGF activity in the un-cleaved form, underscoring that stabilization of the SPH domain is critical for HGF function as a MET ligand [82]. However, the mechanism by which binding of the β -chain to MET sema domain exerts its activating effect on the receptor is unclear. Crystal contacts observed between adjacent SPH domains in the structure of the SPH/MET complex have been proposed as a potential HGF dimerization mechanism (Figure 3F). This observation still awaits experimental validation and as discussed below, it is the α -chain of HGF that likely directly supports MET receptor dimerization.

Insights into HGF α -chain binding and autoinhibition from studies of its fragments: NK1 and NK2

Our understanding of how the HGF α -chain binds to MET and contributes to its dimerization has been predominantly inferred from detailed biochemical and structural analysis of the NK1 and NK2 ligands alone. These two ligands have opposing effects on MET activation. NK1 contains the N domain and the first kringle domain (K1) and is a MET agonist; NK2 contains the N domain and the first two kringle domains (K1 and K2) and is a MET antagonist [83–85] (Figure 1C). Despite the potential mechanistic differences between NK1 and HGF in MET activation discussed below, studies of NK1 and NK2 have provided important insights into how these fragments might behave in the context of the full-length ligand.

The NK1 fragment of the HGF α -chain contains the high-affinity binding site for the MET ECD ($K_d \sim 150$ nM) [77]. Both N and K domains alone can also bind MET, albeit with low affinity ($K_d \sim 1-3 \mu$ M) [77]. NK1 function as a ligand is dependent on heparan sulfate binding and the N domain contains a high-affinity binding site for heparin or heparan sulfate [84,86,87] (Figure 4A). Heparan sulfate induces NK1 dimerization in solution and increases its affinity for MET ($K_d \sim 3$ nM), which is equivalent to the biological activity of the mature HGF molecule in cells [83–85]. A direct MET binding site on N domain is not characterized but we know that K1 domain binds to MET via an interface that is commonly used by kringle domains to bind lysine residues (lysine binding site, LBS) [21,62,88]. The non-canonical LBS of the HGF K1 domain has diverged from other kringle domains and is predicted not to bind lysine side chains [89,90]. While the LBS site is critical for NK1 activity, its precise role in full-length HGF is still contentious. Mutation of LBS site on the K1 domain of HGF abrogates binding to MET [90]. However, a recent study demonstrated that while mutation of the K1 binding site residues in full-length HGF decreased binding of the un-cleaved ligand, in its cleaved (mature) form these mutants could still bind the MET ECD and were equally potent as the wild type, mature HGF in activating MET signaling [91].

Structural analysis of isolated NK1 and NK2 ligands have elucidated the mechanisms by which they act as an agonist and an antagonist of MET, respectively. Crystals structures of the NK1 ligand consistently show a head to tail dimer, with the dimerization interface formed via the linker region between the N and K1 domains as well as reciprocal binding between the N and K1 domains of opposing ligand monomers [87,89,90,92–94] (Figure 4A). In this orientation, the two LBS motifs in the K1 domains are exposed on opposite sides of the NK1 dimer, providing a hypothesis as to how NK1 brings together two MET receptor molecules for activation (Figure 4A). In NK2, the additional kringle domain (K2) forms interdomain contacts with the N domain, obscuring the dimerization interface thus rendering NK2 monomeric and a MET antagonist [85] (Figure 4B). Mutation of the inhibitory interface between the N and K2 domains turns NK2 into a MET agonist [85]. Similar to NK2, the full-length α -chain of HGF (N-K1-K2-K3-K4) is a MET antagonist [95–97], suggesting that it preserves the intermolecular inhibition enforced by the K2 domain. This mechanism to our knowledge has not been investigated in the full-length ligand.

Additional insights into intramolecular inhibition within HGF come from its comparison to plasminogen. In addition to its active SPH domain (β -chain), plasminogen also contains an N domain and five kringle domains (α -chain) (Figure 4D). In crystal structures of its full-length un-cleaved form, plasminogen is in an autoinhibited conformation stabilized in part through intramolecular interactions between lysine residues in the N domain with the LBS region spanning the K4 and K5 domains [98,99]. In this conformation the cleavage site is inaccessible. The binding of plasminogen kringle domains to lysine residues on fibrin is thought to open the molecule and expose the cleavage motif [100] (Figure 4D). While interdomain interactions in plasminogen are not equivalent to those observed in the autoinhibited NK2 structure [23], it is tempting to speculate that HGF binding to MET via its K1 domain serves to initiate a series of similar interdomain rearrangements. Those





Figure 4. Insights into HGF autoinhibition from the structures of NK1 and NK2 ligands.

Part 1 of 2

(A) Domain organization of the head-to-tail dimer of NK1 is shown schematically in the top left corner. Crystal structure of the NK1 dimer is shown in cartoon representation, with the second monomer shown in a semi-transparent surface mode (PDB: 1GMO). Heparan sulfate is shown in yellow bound to each N domain. Residues important for MET binding in the K1 domain are shown in orange in stick representation. The linker region between the N and K1 domains as well as reciprocal interfaces of the N and K1' and K1 and N' domains are shown contributing to the dimer interface. (B) Domain organization of the



Figure 4. Insights into HGF autoinhibition from the structures of NK1 and NK2 ligands.

Part 2 of 2

autoinhibited NK2 monomer is shown schematically in the top left corner. The NK2 crystal structure is shown in cartoon representation (PDB: 3SP8). The heparan sulfate is modeled in via the alignment of the N domain of the NK2 ligand on the N domain of the NK1 ligand (PDB: 1GMO). The obscured N–K1 linker is indicated by an arrow. (**C**) Cartoon schematic of NK1, NK2 and HGF (cleaved and un-cleaved). The relative binding affinity of these different ligand states for MET is indicated by '+' (low affinity) or '++' (high affinity). The ability of these ligand states to activate MET signaling is denoted by 'X' or '**/**'. Heparan sulfate (HS) binding to the N domain has different effects on these ligands. NK1: increased binding affinity for MET and ligand dimerization (++), NK2: no effect, HGF-un-cleaved: no defined contribution (?), HGF-cleaved: HS promotes HGF dimerization (+) but is not required for signaling. The binding site for MET in the K1 domain and the cleaved SPH domain is indicated by a black and yellow star, respectively. (**D**) Cartoon schematic showing domain architecture and activation of plasminogen. The inactive molecule is maintained via multiple autoinhibitory intramolecular interactions. Substrate binding by the exposed LBS of K1 domain leads to elongation and accessibility to the additional LBS sites and of the cleavage motif. Maturation of plasminogen by cleavage activates the serine protease domain (SP).

would lead in turn to the accessibility of the HGF cleavage site prefacing HGF cleavage by its binding to MET. Indeed, analysis of the full HGF molecule by low-resolution cryo-EM, SAXS and high-speed atomic force microscopy points to a more elongated (open) architecture of cleaved HGF compared with its un-cleaved form [21,101]. The cleaved HGF is also more dynamic, which seems to be essential for its activity as a macrocyclic peptide or a monoclonal antibody that bind to surfaces exposed on the cleaved HGF ligand (between K4 and SPH) inhibit HGF by impairing this dynamic motion [101,102].

Understanding the role of heparan sulfate in MET signaling

Heparan sulfate is an ancillary component of the active HGF/MET complex but the precise role it plays in ligand-induced activation of MET is not entirely clear. Apparent differences between the roles of heparan sulfate in HGF- versus NK1-mediated activation of MET exist. While not strictly required for HGF signaling, heparan sulfate potentiates HGF activation of MET in cells and promotes dimerization of full-length HGF [7,24,25]. The shorter NK1 ligand is strictly dependent on heparan sulfate for dimerization and activation of the receptor [83,84] (Figure 4C). Heparan sulfate also enhances the affinity of NK1 for MET, a function which is independent of its ability to dimerize the ligand [94] (Figure 4C). Surprisingly, the mechanisms by which heparan sulfate facilitates ligand dimerization (HGF and NK1) or enhances affinity (NK1) are not known. Crystal structures of apo NK1 and of NK1 bound to heparan sulfate alters the ligand [87].

Similarly to full-length HGF, the NK2 variant binds MET with high affinity independently of heparan sulfate [85] (Figure 4C). This suggests that the K2 domain might stabilize the MET binding interface in the NK2 and full-length HGF, and in the NK1 variant, which misses the K2 domain, heparan sulfate plays this stabilizing role. It is likely that the answer to these mechanistic distinctions will only be provided by the structural characterization of the HGF α -chain bound to MET. Collectively studies of NK1, NK2 and HGF suggest that HGF engages MET via multiple interfaces and that receptor activation likely follows a stepwise model in which HGF cleavage, ligand binding to heparan sulfate, MET binding by the HGF α -chain and β -chain, and receptor dimerization are all in concert to activate the receptor.

Activation and regulation of MET at the membrane and inside the cell

In contrast with substantial structural characterization of the MET ECD and HGF fragments, the intracellular portion of MET remains largely uncharacterized except for the kinase domain (KD). More than seventy deposited crystal structures of the MET kinase domain have revealed that MET contains a canonical KD (Figure 5A,B) [103]. A majority of MET kinase structures include an additional N-terminal α -helix (JM helix, Asn 1058–Val 1070) that packs on top of the kinase N-lobe and is likely required for stability of the kinase fold (Figure 5B,D). In the inactive state, MET kinase adopts a Src/CDK-like inactive conformation in which the activation loop forms a partial helix and packs against the catalytic cleft, displacing helix α C away from the active site and preventing the helix α C Glu 1127 from engaging the catalytic lysine (Lys 1110) (Figure 5B). The activation loop is tethered to the kinase through multiple interactions, including Tyr 1234 engaging the catalytic Glu 1127, Tyr 1230, and Lys 1232 binding the P-loop, and the active site is partially blocked by Met 1229 [104]. Activation of the MET





Figure 5. MET kinase domain states.

(A) Cartoon representation of the MET receptor intracellular domains. The region encoded by exon 14 is colored yellow. Two residues (Ser 985, Tyr 1003) that are important for MET regulation upon being phosphorylated are marked. The α C helix and JM helix in the kinase domain are indicated in blue and pink, respectively. The activation loop is shown in red. (B) Crystal structures of an inactive (PDB: 2G15) and active (PDB: 3R7O) MET kinase domain are shown in cartoon representation. (C, left panel) The activation loop of the inactive MET structure forms a short helix that obscures the α C-helix from swinging into the active site and from forming the salt bridge required for catalysis (Lys 1110 and Glu 1127). In this Src/CDK-like inactive conformation, the DFG motif is in the DFG-in conformation, as indicated by the Asp 1222 pointing towards the active site. The JM helix is shown in light pink. (C, right panel) In the activation loop from the activation loop tyrosines (Tyr 1234 and Tyr 1235) are phosphorylated (denoted as 'pY'), releasing the activation loop from the active site. (D) Zoomed-in view of the hydrophobic residues that form the interface between the JM helix and the α C helix in the MET kinase domain (PDB: 2G15), including Leu 1058, Leu 1062, Val 1066 on the JM helix and Val 1121, Leu 1125, Iso 1129 on the α C-helix. (E) The kinase proximal region of the C-terminal tail (Gly 1346-Lys 1360) is shown in black bound to the back of the kinase C-lobe (PDB: 1R0P). Tyr 1349 and Tyr 1356 are shown in purple.

kinase domain requires sequential phosphorylation of tandem tyrosine residues in the activation loop (Tyr 1235 and Tyr 1234). Tyr 1235 is first phosphorylated due to greater solvent exposure, thereby promoting destabilization of activation loop interactions, followed by phosphorylation of Tyr 1234, which together stabilize the active state of the kinase [104,105] (Figure 5C). Consequently, mutation of Tyr 1234 to a phenylalanine results in reduced kinase activity through stabilization of the autoinhibited state [105,106].

MET cancer mutations, described in papillary renal cell carcinoma (PRCC) and other solid cancers including hepatocellular carcinomas and advanced head and neck squamous cell carcinoma (HNSCC) [12,107,108], act predominantly through destabilization of the autoinhibited kinase structure [103–106,109]. For example, mutation of the activation loop Tyr 1235 to a negatively charged aspartate, found in PRCC, HNSCC and lung adenocarcinoma, destabilizes the autoinhibited conformation of the activation loop. Likewise, Asp 1228 mutations (His/Val/Asn), detected in PRCC, break an active site salt bridge with Lys 1240. Mutations of Tyr 1230 (His/Asp/Cys) that also occur in PRCC, interrupt interactions observed between the hydrophobic P-loop and the activation loop needed to stabilize ATP binding [103–106,109,110].

Much less is known about the structure of other intracellular domains of MET. The juxtamembrane (JM) region in MET is substantially longer (107 residues) in comparison with most other RTKs and is predicted to be largely unstructured. It has two important regulatory motifs located within the membrane-proximal region (defined here as amino acids Asp 963–Asp 1010, encoded by exon 14) (Figure 5A). One of them is centered around Tyr 1003, which when phosphorylated recruits the E3-ligase Cbl leading to MET endocytosis and lyso-somal degradation [111,112]. The second site involves Ser 985, which is phosphorylated by PKC and also results in enhanced MET internalization and down-regulation [113–115]. The JM segment encoded by human exon 14 is skipped in ~4% of non-small-cell lung cancer patients due to mutations in the intron/exon boundary of exon 14 [116–118]. One of the functional consequences of exon 14 skipping is loss of Cbl-mediated ubiquitination and receptor internalization, leading to prolonged HGF-induced signaling [119,120]. Single point mutation of Tyr



1003 to Phe is also transformative in cell-based studies [111,118]. However, there is evidence for other regulatory elements in the exon 14 besides Cbl binding. The addition of the exon 14 coding sequence to the cytoplasmic TPR-MET, in which exon 14 coding region is lost from the MET ICD during the gene fusion event, reduces transforming capacity even when the Cbl-binding site is mutated [120–122]. This suggests that the inhibitory role of the membrane-proximal half of the JM domain might extend beyond the control of receptor ubiquitination-dependent down-regulation and could potentially involve direct interactions with the KD.

On the opposite site of the KD, the C-terminal tail has been directly implicated in the regulation of MET kinase activity. Addition of peptides mimicking the C-terminal tail to cells inhibited MET autophosphorylation and MET-dependent signaling [123]. Consequently, truncation of the entire tail increased receptor activation [124]. A similar observation has been made for the RON receptor, where C-terminal tail truncations also activated the kinase [125]. Structurally, the resolved kinase proximal region of the C-terminal tail wraps around the back side of the C-lobe towards the activation loop [103] (Figure 5E). Further extension of the C-terminal tail may partially block the substrate-binding region or stabilize the activation loop in an inactive conformation, before Tyr 1349 and Tyr 1356 become phosphorylated and disengage from the kinase C-lobe (Figure 5E). Interestingly, in the case of MET, removal of the most terminal 26 amino acids, which correspond to approximately half of the entire C-terminal tail length, results in decreased receptor activation rather than activation [124]. These early studies highlight a complex role that the C-terminal tail plays in the regulation of receptor activity, where in addition to providing the docking sites for downstream effectors, the tail may also play direct positive and negative regulatory roles in regulating kinase activity.

The TM of the receptor might also play an important role since it contains a GxxxG motif (Gly 933–Leu-Iso–Ala–Gly 937) that facilitates homo-dimerization between membrane-embedded helices [126–128]. Peptides that correspond to the MET TM can form homodimers in an *Escherichia coli*-based membrane dimerization assay [129]. How all the intracellular domains come together upon ligand binding to form an active MET receptor complex is largely unknown.

Conclusions

Given the complexity of the HGF and MET structures and their cleavage-dependent rearrangements, it is not surprising that studies of individual domains of these proteins have not yet revealed all the features of receptor activation. While we know that HGF needs to be cleaved to promote the formation of the active MET receptor complex, we do not know all its binding interfaces on MET and how it co-operates with heparin sulfate to promote the formation of active MET dimers. The structure of the full-length HGF that encompasses both α - and β -chains is still missing, and the structure of full-length MET ECD and the importance of its cleavage remains unknown. Consequently, HGF-dependent structural rearrangements within the MET ECD remain uncharacterized. Lastly, we have no knowledge on how the conformational transitions in the MET ECD translate to the intracellular juxtamembrane and kinase domains. The apparent promiscuity of MET in forming co-complexes with other ligands and membrane receptors adds another dimension of mystery to understanding how MET cellular signaling is regulated at the level of receptor activation. While revealing any of these long-standing MET secrets will not be trivial, it is highly anticipated and has the potential to inform new and more effective strategies for disrupting misregulated MET signaling in disease.

Perspectives

- The MET receptor tyrosine kinase and its cognate ligand HGF comprise a signaling axis essential for development, wound healing and tissue homeostasis. Aberrant HGF/MET signaling has been a target of therapies in many human cancers.
- Understanding the mechanisms of HGF-mediated MET activation and signaling is dependent on the characterization of its basal and activated states. Studies on the fragments of HGF and MET proteins support a model in which HGF is activated by cleavage to form a dimer, which then engages two MET monomers to form an active complex. However, currently, there are no high-resolution structures of the active HGF/MET dimer complex that directly visualize these ligand/receptor interactions.



 Defining the ligand-binding site of the HGF α-chain on MET, along with complete structures of the inactive and active, full-length HGF ligand is needed to fully understand how the basal activity of HGF and its binding to MET is regulated.

Competing Interests

E.A.C. is consultant at Takeda, Merck, Loxo and Pear Diagnostics, reports receiving commercial research grants from Astra Zeneca, Ferro Therapeutics, Senti Biosciences, Merck KgA and Bayer and stock ownership of Tatara Therapeutics, Clara Health, BloodQ, Guardant Health. All other authors declare no competing interests. J.S.F is an advisor for, holds equity in, and has received sponsored research from Relay Therapeutics. N.J. is a member of the SAB and a shareholder of Turning Point Therapeutics, SUDO Biosciences and Type6 Therapeutics. The Jura laboratory has received sponsored research support from Genentech.

Author Contributions

E.M.L., G.O.E. and N.J. wrote the manuscript and designed figures. All authors were involved in the editing and discussion of the manuscript.

Acknowledgements

We thank Raphael Trenker and Devan Diwanji of the Jura lab for ongoing and insightful discussions regarding the regulation of MET and HGF. The authors sincerely apologize to all colleagues whose work was omitted in this review owing to space constraints. This work was supported in part by grants from the National Cancer Institute to N.J. and E.A.C. (R01 CA230263) and E.A.C. (R01 CA239604, R01 CA222862, R01 CA227807, U24 CA210974).

Abbreviation

ECD, extracellular domain; EM, electron microscopy; HGF, hepatocyte growth factor; HGFR, hepatocyte growth factor receptor; HNSCC, head and neck squamous cell carcinoma; IPT, integrin/plexin/transcription; JM, juxtamembrane; KD, kinase domain; LRR, leucine-rich repeat; PRCC, papillary renal cell carcinoma; PSI, plexin/ semaphorin/integrin; RON, recepteur d'Origine nantais; RTK, receptor tyrosine kinase; SAXS, small-angle X-ray scattering; SPH, serine protease homology; TM, transmembrane

References

- 1 Cooper, C.S., Park, M., Blair, D.G., Tainsky, M.A., Huebner, K., Croce, C.M. et al. (1984) Molecular cloning of a new transforming gene from a chemically transformed human cell line. *Nature* **311**, 29–33 https://doi.org/10.1038/311029a0
- 2 Park, M., Dean, M., Kaul, K., Braun, M.J., Gonda, M.A. and Woude G, V. (1987) Sequence of MET protooncogene cDNA has features characteristic of the tyrosine kinase family of growth-factor receptors. *Proc. Natl Acad. Sci. U.S.A.* 84, 6379–6383 https://doi.org/10.1073/pnas.84.18.6379
- Bottaro, D.P., Rubin, J.S., Faletto, D.L., Chan, A.M., Kmiecik, T.E., Vande Woude, G.F. et al. (1991) Identification of the hepatocyte growth factor receptor as the c-met proto-oncogene product. *Science* **251**, 802–804 https://doi.org/10.1126/science.1846706
- 4 Ronsin, C., Muscatelli, F., Mattei, M.G. and Breathnach, R. (1993) A novel putative receptor protein tyrosine kinase of the met family. *Oncogene* **8**, 1195–1202 PMID: 8386824
- 5 Lemmon, M.A. and Schlessinger, J. (2010) Cell signaling by receptor tyrosine kinases. *Cell* **141**, 1117–1134 https://doi.org/10.1016/j.cell.2010.06. 011
- 6 Bork, P., Doerks, T., Springer, T.A. and Snel, B. (1999) Domains in plexins: links to integrins and transcription factors. *Trends Biochem. Sci.* 24, 261–263 https://doi.org/10.1016/S0968-0004(99)01416-4
- 7 Gherardi, E., Youles, M.E., Miguel, R.N., Blundell, T.L., lamele, L., Gough, J. et al. (2003) Functional map and domain structure of MET, the product of the c-met protooncogene and receptor for hepatocyte growth factor/scatter factor. *Proc. Natl Acad. Sci. U.S.A.* **100**, 12039–12044 https://doi.org/10. 1073/pnas.2034936100
- 8 Trusolino, L., Bertotti, A. and Comoglio, P.M. (2010) MET signalling: principles and functions in development, organ regeneration and cancer. Nat. Rev. Mol. Cell Biol. 11, 834–848 https://doi.org/10.1038/nrm3012
- 9 Petrini, I. (2015) Biology of MET: a double life between normal tissue repair and tumor progression. Ann. Transl. Med. 3, 82 https://doi.org/10.3978/j. issn.2305-5839.2015.03.58
- 10 Kato, T. (2017) Biological roles of hepatocyte growth factor-Met signaling from genetically modified animals (Review). *Biomed. Rep.* **7**, 495–503 https://doi.org/10.3892/br.2017.1001
- 11 De Silva, D.M., Roy, A., Kato, T., Cecchi, F., Lee, Y.H., Matsumoto, K. et al. (2017) Targeting the hepatocyte growth factor/Met pathway in cancer. Biochem. Soc. Trans. 45, 855–870 https://doi.org/10.1042/BST20160132



- 12 Comoglio, P.M., Trusolino, L. and Boccaccio, C. (2018) Known and novel roles of the MET oncogene in cancer: a coherent approach to targeted therapy. *Nat. Rev. Cancer* **18**, 341–358 https://doi.org/10.1038/s41568-018-0002-y
- 13 Koch, J.P., Aebersold, D.M., Zimmer, Y. and Medová, M. (2020) MET targeting: time for a rematch. *Oncogene* **39**, 2845–2862 https://doi.org/10. 1038/s41388-020-1193-8
- 14 Nakamura, T., Nawa, K., Ichihara, A., Kaise, N. and Nishino, T. (1987) Purification and subunit structure of hepatocyte growth factor from rat platelets. *FEBS Lett.* **224**, 311–316 https://doi.org/10.1016/0014-5793(87)80475-1
- 15 Stoker, M., Gherardi, E., Perryman, M. and Gray, J. (1987) Scatter factor is a fibroblast-derived modulator of epithelial cell mobility. *Nature* **327**, 239–242 https://doi.org/10.1038/327239a0
- 16 Gherardi, E. and Stoker, M. (1990) Hepatocytes and scatter factor. Nature 346, 228 https://doi.org/10.1038/346228b0
- 17 Weidner, K.M., Arakaki, N., Hartmann, G., Vandekerckhove, J., Weingart, S., Rieder, H. et al. (1991) Evidence for the identity of human scatter factor and human hepatocyte growth factor. *Proc. Natl Acad. Sci. U.S.A.* **88**, 7001–705 https://doi.org/10.1073/pnas.88.16.7001
- 18 Nakamura, T., Nishizawa, T., Hagiya, M., Seki, T., Shimonishi, M., Sugimura, A. et al. (1989) Molecular cloning and expression of human hepatocyte growth factor. *Nature* **342**, 440–443 https://doi.org/10.1038/342440a0
- 19 Cioce, V., Csaky, K.G., Chan, A.M., Bottaro, D.P., Taylor, W.G., Jensen, R. et al. (1996) Hepatocyte growth factor (HGF)/NK1 is a naturally occurring HGF/scatter factor variant with partial agonist/antagonist activity. *J. Biol. Chem.* **271**, 13110–13115 https://doi.org/10.1074/jbc.271.22. 13110
- 20 Chan, A.M., Rubin, J.S., Bottaro, D.P., Hirschfield, D.W., Chedid, M. and Aaronson, S.A. (1991) Identification of a competitive HGF antagonist encoded by an alternative transcript. *Science* **254**, 1382–1385 https://doi.org/10.1126/science.1720571
- 21 Gherardi, E., Sandin, S., Petoukhov, M.V., Finch, J., Youles, M.E., Ofverstedt, L.G. et al. (2006) Structural basis of hepatocyte growth factor/scatter factor and MET signalling. *Proc. Natl Acad. Sci. U.S.A.* **103**, 4046–4051 https://doi.org/10.1073/pnas.0509040103
- 22 Gherardi, E., Birchmeier, W., Birchmeier, C. and Vande Woude, G. (2012) Targeting MET in cancer: rationale and progress. Nat. Rev. Cancer 12, 89–103 https://doi.org/10.1038/nrc3205
- 23 Niemann, H.H. (2013) Structural basis of MET receptor dimerization by the bacterial invasion protein inIB and the HGF/SF splice variant NK1. *Biochim. Biophys. Acta* 1834, 2195–2204 https://doi.org/10.1016/j.bbapap.2012.10.012
- 24 Zioncheck, T.F., Richardson, L., Liu, J., Chang, L., King, K.L., Bennett, G.L. et al. (1995) Sulfated oligosaccharides promote hepatocyte growth factor association and govern its mitogenic activity. J. Biol. Chem. 270, 16871–16878 https://doi.org/10.1074/jbc.270.28.16871
- 25 Kemp, L.E., Mulloy, B. and Gherardi, E. (2006) Signalling by HGF/SF and Met: the role of heparan sulphate co-receptors. *Biochem. Soc. Trans.* 34(Pt 3), 414–417 https://doi.org/10.1042/BST0340414
- 26 Shen, Y., Naujokas, M., Park, M. and Ireton, K. (2000) InIB-dependent internalization of Listeria is mediated by the Met receptor tyrosine kinase. *Cell* 103, 501–510 https://doi.org/10.1016/S0092-8674(00)00141-0
- 27 Niemann, H.H., Jager, V., Butler, P.J., van den Heuvel, J., Schmidt, S., Ferraris, D. et al. (2007) Structure of the human receptor tyrosine kinase met in complex with the Listeria invasion protein InIB. *Cell* **130**, 235–246 https://doi.org/10.1016/j.cell.2007.05.037
- 28 Radoshevich, L. and Cossart, P. (2018) *Listeria monocytogenes*: towards a complete picture of its physiology and pathogenesis. *Nat. Rev. Microbiol.* **16**, 32–46 https://doi.org/10.1038/nrmicro.2017.126
- 29 Goldoni, S., Humphries, A., Nyström, A., Sattar, S., Owens, R.T., McQuillan, D.J. et al. (2009) Decorin is a novel antagonistic ligand of the Met receptor. J. Cell Biol. **185**, 743–754 https://doi.org/10.1083/jcb.200901129
- 30 Trusolino, L., Bertotti, A. and Comoglio, P.M. (2001) A signaling adapter function for alpha6beta4 integrin in the control of HGF-dependent invasive growth. Cell 107, 643–654 https://doi.org/10.1016/S0092-8674(01)00567-0
- 31 Orian-Rousseau, V., Chen, L., Sleeman, J.P., Herrlich, P. and Ponta, H. (2002) CD44 is required for two consecutive steps in HGF/c-Met signaling. *Genes Dev.* **16**, 3074–3086 https://doi.org/10.1101/gad.242602
- 32 Lu, K.V., Chang, J.P., Parachoniak, C.A., Pandika, M.M., Aghi, M.K., Meyronet, D. et al. (2012) VEGF inhibits tumor cell invasion and mesenchymal transition through a MET/VEGFR2 complex. *Cancer Cell* 22, 21–35 https://doi.org/10.1016/j.ccr.2012.05.037
- 33 Gentile, A., Lazzari, L., Benvenuti, S., Trusolino, L. and Comoglio, P.M. (2014) The ROR1 pseudokinase diversifies signaling outputs in MET-addicted cancer cells. *Int. J. Cancer* **135**, 2305–2316 https://doi.org/10.1002/ijc.28879
- 34 Frazier, N.M., Brand, T., Gordan, J.D., Grandis, J. and Jura, N. (2019) Overexpression-mediated activation of MET in the Golgi promotes HER3/ERBB3 phosphorylation. *Oncogene* 38, 1936–1950 https://doi.org/10.1038/s41388-018-0537-0
- 35 Kong, L.R., Mohamed Salleh, N.A.B., Ong, R.W., Tan, T.Z., Syn, N.L., Goh, R.M. et al. (2020) A common MET polymorphism harnesses HER2 signaling to drive aggressive squamous cell carcinoma. *Nat. Commun.* **11**, 1556 https://doi.org/10.1038/s41467-020-15318-5
- 36 Li, W., Xiong, X., Abdalla, A., Alejo, S., Zhu, L., Lu, F. et al. (2018) HGF-induced formation of the MET-AXL-ELM02-DOCK180 complex promotes RAC1 activation, receptor clustering, and cancer cell migration and invasion. J. Biol. Chem. 293, 15397–15418 https://doi.org/10.1074/jbc.RA118.003063
- 37 Benvenuti, S., Lazzari, L., Arnesano, A., Li Chiavi, G., Gentile, A. and Comoglio, P.M. (2011) Ron kinase transphosphorylation sustains MET oncogene addiction. *Cancer Res.* **71**, 1945–1955 https://doi.org/10.1158/0008-5472.CAN-10-2100
- 38 Giordano, S., Corso, S., Conrotto, P., Artigiani, S., Gilestro, G., Barberis, D. et al. (2002) The semaphorin 4D receptor controls invasive growth by coupling with Met. Nat. Cell Biol. 4, 720–724 https://doi.org/10.1038/ncb843
- 39 Lai, A.Z., Abella, J.V. and Park, M. (2009) Crosstalk in Met receptor oncogenesis. *Trends Cell Biol.* **19**, 542–551 https://doi.org/10.1016/j.tcb.2009.07.002
- 40 Viticchie, G. and Muller, P.A.J. (2015) c-Met and other cell surface molecules: interaction. *Act. Funct. Conseq. Biomed.* **3**, 46–70 https://doi.org/10. 3390/biomedicines3010046
- 41 Bardelli, A., Corso, S., Bertotti, A., Hobor, S., Valtorta, E., Siravegna, G. et al. (2013) Amplification of the MET receptor drives resistance to anti-EGFR therapies in colorectal cancer. *Cancer Discov.* 3, 658–673 https://doi.org/10.1158/2159-8290.CD-12-0558
- 42 Diwanji, D., Thaker, T. and Jura, N. (2019) More than the sum of the parts: Toward full-length receptor tyrosine kinase structures. *IUBMB Life* **71**, 706–720 https://doi.org/10.1002/iub.2060
- 43 Gherardi, E., Love, C.A., Esnouf, R.M. and Jones, E.Y. (2004) The sema domain. Curr. Opin. Struct. Biol. 14, 669–678 https://doi.org/10.1016/j.sbi.2004.10.010
- 44 Pascoe, H.G., Wang, Y. and Zhang, X. (2015) Structural mechanisms of plexin signaling. *Prog. Biophys. Mol. Biol.* **118**, 161–168 https://doi.org/10. 1016/j.pbiomolbio.2015.03.006



- 45 Alves, J.M., Martins, A.H., Lameu, C., Glaser, T., Boukli, N.M., Bassaneze, V. et al. (2019) Kinin-B2 receptor activity in skeletal muscle regeneration and myoblast differentiation. *Stem Cell Rev. Rep.* **15**, 48–58 https://doi.org/10.1007/s12015-018-9850-9
- 46 Siebold, C. and Jones, E.Y. (2013) Structural insights into semaphorins and their receptors. *Semin. Cell Dev. Biol.* 24, 139–145 https://doi.org/10. 1016/j.semcdb.2012.11.003
- 47 Love, C.A., Harlos, K., Mavaddat, N., Davis, S.J., Stuart, D.I., Jones, E.Y. et al. (2003) The ligand-binding face of the semaphorins revealed by the high-resolution crystal structure of SEMA4D. *Nat. Struct. Biol.* **10**, 843–848 https://doi.org/10.1038/nsb977
- 48 Antipenko, A., Himanen, J.P., van Leyen, K., Nardi-Dei, V., Lesniak, J., Barton, W.A. et al. (2003) Structure of the semaphorin-3A receptor binding module. *Neuron* **39**, 589–598 https://doi.org/10.1016/S0896-6273(03)00502-6
- 49 Janssen, B.J., Robinson, R.A., Perez-Branguli, F., Bell, C.H., Mitchell, K.J., Siebold, C. et al. (2010) Structural basis of semaphorin-plexin signalling. *Nature* 467, 1118–1122 https://doi.org/10.1038/nature09468
- 50 Liu, H., Juo, Z.S., Shim, A.H., Focia, P.J., Chen, X., Garcia, K.C. et al. (2010) Structural basis of semaphorin-plexin recognition and viral mimicry from Sema7A and A39R complexes with PlexinC1. *Cell* **142**, 749–761 https://doi.org/10.1016/j.cell.2010.07.040
- 51 Nogi, T., Yasui, N., Mihara, E., Matsunaga, Y., Noda, M., Yamashita, N. et al. (2010) Structural basis for semaphorin signalling through the plexin receptor. *Nature* **467**, 1123–1127 https://doi.org/10.1038/nature09473
- 52 Stamos, J., Lazarus, R.A., Yao, X., Kirchhofer, D. and Wiesmann, C. (2004) Crystal structure of the HGF beta-chain in complex with the sema domain of the Met receptor. *EMBO J.* 23, 2325–2335 https://doi.org/10.1038/sj.emboj.7600243
- 53 Komada, M., Hatsuzawa, K., Shibamoto, S., Ito, F., Nakayama, K. and Kitamura, N. (1993) Proteolytic processing of the hepatocyte growth factor/scatter factor receptor by furin. *FEBS Lett.* **328**, 25–29 https://doi.org/10.1016/0014-5793(93)80958-W
- 54 Mark, M.R., Lokker, N.A., Zioncheck, T.F., Luis, E.A. and Godowski, P.J. (1992) Expression and characterization of hepatocyte growth factor receptor-IgG fusion proteins. Effects of mutations in the potential proteolytic cleavage site on processing and ligand binding. *J. Biol. Chem.* **267**, 26166–26171 https://doi.org/10.1016/S0021-9258(18)35731-4
- 55 Niemann, H.H. (2011) Structural insights into Met receptor activation. Eur. J. Cell Biol. 90, 972–981 https://doi.org/10.1016/j.ejcb.2010.11.014
- 56 Chao, K.L., Tsai, I.W., Chen, C. and Herzberg, O. (2012) Crystal structure of the sema-PSI extracellular domain of human RON receptor tyrosine kinase. PLoS One 7, e41912 https://doi.org/10.1371/journal.pone.0041912
- 57 DiCara, D.M., Chirgadze, D.Y., Pope, A.R., Karatt-Vellatt, A., Winter, A., Slavny, P. et al. (2017) Characterization and structural determination of a new anti-MET function-blocking antibody with binding epitope distinct from the ligand binding domain. *Sci. Rep.* **7**, 9000 https://doi.org/10.1038/ s41598-017-09460-2
- 58 Niemann, H.H., Petoukhov, M.V., Hartlein, M., Moulin, M., Gherardi, E., Timmins, P. et al. (2008) X-ray and neutron small-angle scattering analysis of the complex formed by the Met receptor and the *Listeria monocytogenes* invasion protein InIB. *J. Mol. Biol.* **377**, 489–500 https://doi.org/10.1016/j.jmb.2008.01.027
- 59 Kong, Y., Janssen, B.J., Malinauskas, T., Vangoor, V.R., Coles, C.H., Kaufmann, R. et al. (2016) Structural basis for plexin activation and regulation. *Neuron* **91**, 548–560 https://doi.org/10.1016/j.neuron.2016.06.018
- 60 Suzuki, K., Tsunoda, H., Omiya, R., Matoba, K., Baba, T., Suzuki, S. et al. (2016) Structure of the plexin ectodomain bound by Semaphorin-Mimicking antibodies. *PLoS One* **11**, e0156719. https://doi.org/10.1371/journal.pone.0156719
- 61 Kong-Beltran, M., Stamos, J. and Wickramasinghe, D. (2004) The sema domain of Met is necessary for receptor dimerization and activation. *Cancer Cell* **6**, 75–84 https://doi.org/10.1016/j.ccr.2004.06.013
- 62 Youles, M., Holmes, O., Petoukhov, M.V., Nessen, M.A., Stivala, S., Svergun, D.I. et al. (2008) Engineering the NK1 fragment of hepatocyte growth factor/scatter factor as a MET receptor antagonist. J. Mol. Biol. **377**, 616–622 https://doi.org/10.1016/j.jmb.2008.01.034
- 63 Koschut, D., Richert, L., Pace, G., Niemann, H.H., Mély, Y. and Orian-Rousseau, V. (2016) Live cell imaging shows hepatocyte growth factor-induced Met dimerization. *Biochim. Biophys. Acta* 1863, 1552–1558 https://doi.org/10.1016/j.bbamcr.2016.04.015
- 64 Lemmon, M.A., Pinchasi, D., Zhou, M., Lax, I. and Schlessinger, J. (1997) Kit receptor dimerization is driven by bivalent binding of stem cell factor. J. Biol. Chem. 272, 6311–6317 https://doi.org/10.1074/jbc.272.10.6311
- Reshetnyak, A.V., Opatowsky, Y., Boggon, T.J., Folta-Stogniew, E., Tome, F., Lax, I. et al. (2015) The strength and cooperativity of KIT ectodomain contacts determine normal ligand-dependent stimulation or oncogenic activation in cancer. *Mol. Cell* 57, 191–201 https://doi.org/10.1016/j.molcel. 2014.11.021
- 66 Dawson, J.P., Bu, Z. and Lemmon, M.A. (2007) Ligand-induced structural transitions in ErbB receptor extracellular domains. *Structure* **15**, 942–954 https://doi.org/10.1016/j.str.2007.06.013
- 67 Andres, F., Iamele, L., Meyer, T., Stüber, J.C., Kast, F., Gherardi, E. et al. (2019) Inhibition of the MET kinase activity and cell growth in MET-Addicted cancer cells by bi-paratopic linking. J. Mol. Biol. 431, 2020–2039 https://doi.org/10.1016/j.jmb.2019.03.024
- Banerjee, M., Copp, J., Vuga, D., Marino, M., Chapman, T., van der Geer, P. et al. (2004) GW domains of the *Listeria monocytogenes* invasion protein InIB are required for potentiation of Met activation. *Mol. Microbiol.* **52**, 257–271 https://doi.org/10.1111/j.1365-2958.2003.03968.x
- 69 Ferraris, D.M., Gherardi, E., Di, Y., Heinz, D.W. and Niemann, H.H. (2010) Ligand-mediated dimerization of the Met receptor tyrosine kinase by the bacterial invasion protein InIB. *J. Mol. Biol.* **395**, 522–532 https://doi.org/10.1016/j.jmb.2009.10.074
- 70 Basilico, C., Hultberg, A., Blanchetot, C., de Jonge, N., Festjens, E., Hanssens, V. et al. (2014) Four individually druggable MET hotspots mediate HGF-driven tumor progression. *J. Clin. Invest.* **124**, 3172–3186 https://doi.org/10.1172/JCI72316
- 71 Hartmann, G., Naldini, L., Weidner, K.M., Sachs, M., Vigna, E., Comoglio, P.M. et al. (1992) A functional domain in the heavy chain of scatter factor/ hepatocyte growth factor binds the c-Met receptor and induces cell dissociation but not mitogenesis. *Proc. Natl Acad. Sci. U.S.A.* 89, 11574–11578 https://doi.org/10.1073/pnas.89.23.11574
- 72 Lokker, N.A., Mark, M.R., Luis, E.A., Bennett, G.L., Robbins, K.A., Baker, J.B. et al. (1992) Structure-function analysis of hepatocyte growth factor: identification of variants that lack mitogenic activity yet retain high affinity receptor binding. *EMBO J.* **11**, 2503–2510 https://doi.org/10.1002/j. 1460-2075.1992.tb05315.x
- 73 Maun, H.R., Kirchhofer, D. and Lazarus, R.A. (2010) Pseudo-active sites of protease domains: HGF/Met and sonic hedgehog signaling in cancer. *Biol. Chem.* 391881–391892 https://doi.org/10.1515/BC.2010.098
- 74 Naldini, L., Tamagnone, L., Vigna, E., Sachs, M., Hartmann, G., Birchmeier, W. et al. (1992) Extracellular proteolytic cleavage by urokinase is required for activation of hepatocyte growth factor/scatter factor. *EMBO J.* **11**, 4825–4833 https://doi.org/10.1002/j.1460-2075.1992.tb05588.x



- 75 Kirchhofer, D., Yao, X., Peek, M., Eigenbrot, C., Lipari, M.T., Billeci, K.L. et al. (2004) Structural and functional basis of the serine protease-like hepatocyte growth factor β-chain in met binding and signaling. *J. Biol. Chem.* **279**, 39915–39924 https://doi.org/10.1074/jbc. M404795200
- 76 Matsumoto, K., Kataoka, H., Date, K. and Nakamura, T. (1998) Cooperative interaction between alpha- and beta-chains of hepatocyte growth factor on c-Met receptor confers ligand-induced receptor tyrosine phosphorylation and multiple biological responses. J. Biol. Chem. 273, 22913–22920 https://doi.org/10.1074/jbc.273.36.22913
- 77 Holmes, O., Pillozzi, S., Deakin, J.A., Carafoli, F., Kemp, L., Butler, P.J.G. et al. (2007) Insights into the structure/function of hepatocyte growth factor/ scatter factor from studies with individual domains. J. Mol. Biol. 367, 395–408 https://doi.org/10.1016/j.jmb.2006.12.061
- 78 Merchant, M., Ma, X., Maun, H.R., Zheng, Z., Peng, J., Romero, M. et al. (2013) Monovalent antibody design and mechanism of action of onartuzumab, a MET antagonist with anti-tumor activity as a therapeutic agent. *Proc. Natl Acad. Sci. U.S.A.* **110**, E2987–E2996 https://doi.org/10. 1073/pnas.1302725110
- 79 Miao, W., Sakai, K., Sato, H., Imamura, R., Jangphattananont, N., Takagi, J. et al. (2019) Impaired ligand-dependent MET activation caused by an extracellular SEMA domain missense mutation in lung cancer. *Cancer Sci.* **110**, 3340–3349 https://doi.org/10.1111/cas.14142
- 80 Tode, N., Kikuchi, T., Sakakibara, T., Hirano, T., Inoue, A., Ohkouchi, S. et al. (2017) Exome sequencing deciphers a germline MET mutation in familial epidermal growth factor receptor-mutant lung cancer. *Cancer Sci.* **108**, 1263–1270 https://doi.org/10.1111/cas.13233
- 81 Basilico, C., Arnesano, A., Galluzzo, M., Comoglio, P.M. and Michieli, P. (2008) A high affinity hepatocyte growth factor-binding site in the immunoglobulin-like region of Met. *J. Biol. Chem.* **283**, 21267–21277 https://doi.org/10.1074/jbc.M800727200
- 82 Landgraf, K.E., Steffek, M., Quan, C., Tom, J., Yu, C., Santell, L. et al. (2014) An allosteric switch for pro-HGF/Met signaling using zymogen activator peptides. *Nat. Chem. Biol.* **10**, 567–573 https://doi.org/10.1038/nchembio.1533
- 83 Schwall, R.H., Chang, L.Y., Godowski, P.J., Kahn, D.W., Hillan, K.J., Bauer, K.D. et al. (1996) Heparin induces dimerization and confers proliferative activity onto the hepatocyte growth factor antagonists NK1 and NK2. *J. Cell Biol.* **133**, 709–718 https://doi.org/10.1083/jcb.133.3.709
- 84 Sakata, H., Stahl, S.J., Taylor, W.G., Rosenberg, J.M., Sakaguchi, K., Wingfield, P.T. et al. (1997) Heparin binding and oligomerization of hepatocyte growth factor/scatter factor isoforms. Heparan sulfate glycosaminoglycan requirement for Met binding and signaling. J. Biol. Chem. 272, 9457–9463 https://doi.org/10.1074/jbc.272.14.9457
- 85 Tolbert, W.D., Daugherty-Holtrop, J., Gherardi, E., Vande Woude, G. and Xu, H.E. (2010) Structural basis for agonism and antagonism of hepatocyte growth factor. *Proc. Natl Acad. Sci. U.S.A.* **107**, 13264–13269 https://doi.org/10.1073/pnas.1005183107
- 27 Zhou, H., Mazzulla, M.J., Kaufman, J.D., Stahl, S.J., Wingfield, P.T., Rubin, J.S. et al. (1998) The solution structure of the N-terminal domain of hepatocyte growth factor reveals a potential heparin-binding site. *Structure* 6, 109–116 https://doi.org/10.1016/S0969-2126(98) 00012-4
- 87 Lietha, D., Chirgadze, D.Y., Mulloy, B., Blundell, T.L. and Gherardi, E. (2001) Crystal structures of NK1-heparin complexes reveal the basis for NK1 activity and enable engineering of potent agonists of the MET receptor. *EMBO J.* 20, 5543–5555 https://doi.org/10.1093/emboj/20.20.5543
- 88 Lokker, N.A., Presta, L.G. and Godowski, P.J. (1994) Mutational analysis and molecular modeling of the N-terminal kringle-containing domain of hepatocyte growth factor identifies amino acid side chains important for interaction with the c-Met receptor. *Protein Eng.* **7**, 895–903 https://doi.org/10. 1093/protein/7.7.895
- 89 Ultsch, M., Lokker, N.A., Godowski, P.J. and de Vos, A.M. (1998) Crystal structure of the NK1 fragment of human hepatocyte growth factor at 2.0 A resolution. *Structure* 6, 1383–1393 https://doi.org/10.1016/S0969-2126(98)00138-5
- 90 Sigurdardottir, A.G., Winter, A., Sobkowicz, A., Fragai, M., Chirgadze, D., Ascher, D.B. et al. (2015) Exploring the chemical space of the lysine-binding pocket of the first kringle domain of hepatocyte growth factor/scatter factor (HGF/SF) yields a new class of inhibitors of HGF/SF-MET binding. *Chem. Sci.* 6, 6147–6157 https://doi.org/10.1039/C5SC02155C
- 91 Umitsu, M., Sakai, K., Tamura-Kawakami, K., Matsumoto, K. and Takagi, J. (2020) The constitutive high-affinity Met-binding site in the kringle domain is dispensable for the signalling activity of hepatocyte growth factor. *J. Biochem.* **167**, 577–586 https://doi.org/10.1093/jb/mvaa006
- 92 Chirgadze, D.Y., Hepple, J.P., Zhou, H., Byrd, R.A., Blundell, T.L. and Gherardi, E. (1999) Crystal structure of the NK1 fragment of HGF/SF suggests a novel mode for growth factor dimerization and receptor binding. *Nat. Struct. Biol.* **6**, 72–79 https://doi.org/10.1038/4947
- 93 Watanabe, K., Chirgadze, D.Y., Lietha, D., de Jonge, H., Blundell, T.L. and Gherardi, E. (2002) A new crystal form of the NK1 splice variant of HGF/SF demonstrates extensive hinge movement and suggests that the NK1 dimer originates by domain swapping. J. Mol. Biol. 319, 283–288 https://doi.org/ 10.1016/S0022-2836(02)00199-7
- 94 Tolbert, W.D., Daugherty, J., Gao, C., Xie, Q., Miranti, C., Gherardi, E. et al. (2007) A mechanistic basis for converting a receptor tyrosine kinase agonist to an antagonist. Proc. Natl Acad. Sci. U.S.A. 104, 14592–14597 https://doi.org/10.1073/pnas.0704290104
- 95 Date, K., Matsumoto, K., Shimura, H., Tanaka, M. and Nakamura, T. (1997) HGF/NK4 is a specific antagonist for pleiotrophic actions of hepatocyte growth factor. FEBS Lett. 420, 1–6 https://doi.org/10.1016/S0014-5793(97)01475-0
- 96 Date, K., Matsumoto, K., Kuba, K., Shimura, H., Tanaka, M. and Nakamura, T. (1998) Inhibition of tumor growth and invasion by a four-kringle antagonist (HGF/NK4) for hepatocyte growth factor. *Oncogene* 17, 3045–3054 https://doi.org/10.1038/sj.onc.1202231
- 97 Matsumoto, K. and Nakamura, T. (2005) Mechanisms and significance of bifunctional NK4 in cancer treatment. *Biochem. Biophys. Res. Commun.* 333, 316–327 https://doi.org/10.1016/j.bbrc.2005.05.131
- 98 Law, R.H., Caradoc-Davies, T., Cowieson, N., Horvath, A.J., Quek, A.J., Encarnacao, J.A. et al. (2012) The X-ray crystal structure of full-length human plasminogen. *Cell Rep.* 1, 185–190 https://doi.org/10.1016/j.celrep.2012.02.012
- 99 Xue, Y., Bodin, C. and Olsson, K. (2012) Crystal structure of the native plasminogen reveals an activation-resistant compact conformation. *J. Thromb. Haemost.* **10**, 1385–1396 https://doi.org/10.1111/j.1538-7836.2012.04765.x
- 100 Wu, G., Quek, A.J., Caradoc-Davies, T.T., Ekkel, S.M., Mazzitelli, B., Whisstock, J.C. et al. (2019) Structural studies of plasmin inhibition. *Biochem. Soc* . *Trans.* 47, 541–557 https://doi.org/10.1042/BST20180211
- 101 Sakai, K., Passioura, T., Sato, H., Ito, K., Furuhashi, H., Umitsu, M. et al. (2019) Macrocyclic peptide-based inhibition and imaging of hepatocyte growth factor. *Nat. Chem. Biol.* **15**, 598–606 https://doi.org/10.1038/s41589-019-0285-7
- 102 Umitsu, M., Sakai, K., Ogasawara, S., Kaneko, M.K., Asaki, R., Tamura-Kawakami, K. et al. (2016) Probing conformational and functional states of human hepatocyte growth factor by a panel of monoclonal antibodies. *Sci. Rep.* **6**, 33149. https://doi.org/10.1038/srep33149



- 103 Schiering, N., Knapp, S., Marconi, M., Flocco, M.M., Cui, J., Perego, R. et al. (2003) Crystal structure of the tyrosine kinase domain of the hepatocyte growth factor receptor c-Met and its complex with the microbial alkaloid K-252a. *Proc. Natl Acad. Sci. U.S.A.* **100**, 12654–12659 https://doi.org/10. 1073/pnas.1734128100
- 104 Wang, W., Marimuthu, A., Tsai, J., Kumar, A., Krupka, H.I., Zhang, C. et al. (2006) Structural characterization of autoinhibited c-Met kinase produced by coexpression in bacteria with phosphatase. *Proc. Natl Acad. Sci. U.S.A.* **103**, 3563–3568 https://doi.org/10.1073/pnas.0600048103
- 105 Chiara, F., Michieli, P., Pugliese, L. and Comoglio, P.M. (2003) Mutations in the met oncogene unveil a "dual switch" mechanism controlling tyrosine kinase activity. *J. Biol. Chem.* **278**, 29352–29358 https://doi.org/10.1074/jbc.M302404200
- 106 Cristiani, C., Rusconi, L., Perego, R., Schiering, N., Kalisz, H.M., Knapp, S. et al. (2005) Regulation of the wild-type and Y1235D mutant Met kinase activation. *Biochemistry* 44, 14110–14119 https://doi.org/10.1021/bi051242k
- 107 Duplaquet, L., Kherrouche, Z., Baldacci, S., Jamme, P., Cortot, A.B., Copin, M.C. et al. (2018) The multiple paths towards MET receptor addiction in cancer. *Oncogene* **37**, 3200–3215 https://doi.org/10.1038/s41388-018-0185-4
- 108 Recondo, G., Che, J., Janne, P.A. and Awad, M.M. (2020) Targeting MET dysregulation in cancer. *Cancer Discov.* **10**, 922–934 https://doi.org/10. 1158/2159-8290.CD-19-1446
- 109 Timofeevski, S.L., McTigue, M.A., Ryan, K., Cui, J., Zou, H.Y., Zhu, J.X. et al. (2009) Enzymatic characterization of c-Met receptor tyrosine kinase oncogenic mutants and kinetic studies with aminopyridine and triazolopyrazine inhibitors. *Biochemistry* 48, 5339–5349 https://doi.org/10.1021/ bi900438w
- 110 Schmidt, L., Duh, F.M., Chen, F., Kishida, T., Glenn, G., Choyke, P. et al. (1997) Germline and somatic mutations in the tyrosine kinase domain of the MET proto-oncogene in papillary renal carcinomas. *Nat. Genet.* **16**, 68–73 https://doi.org/10.1038/ng0597-68
- 111 Peschard, P., Fournier, T.M., Lamorte, L., Naujokas, M.A., Band, H., Langdon, W.Y. et al. (2001) Mutation of the c-Cbl TKB domain binding site on the Met receptor tyrosine kinase converts it into a transforming protein. *Mol. Cell* **8**, 995–1004 https://doi.org/10.1016/S1097-2765(01)00378-1
- 112 Petrelli, A., Gilestro, G.F., Lanzardo, S., Comoglio, P.M., Migone, N. and Giordano, S. (2002) The endophilin-CIN85-Cbl complex mediates ligand-dependent downregulation of c-Met. *Nature* **416**, 187–190 https://doi.org/10.1038/416187a
- 113 Kermorgant, S., Zicha, D. and Parker, P.J. (2003) Protein kinase C controls microtubule-based traffic but not proteasomal degradation of c-Met. *J. Biol. Chem.* 278, 28921–28929 https://doi.org/10.1074/jbc.M302116200
- 114 Hashigasako, A., Machide, M., Nakamura, T., Matsumoto, K. and Nakamura, T. (2004) Bi-directional regulation of Ser-985 phosphorylation of c-met via protein kinase C and protein phosphatase 2A involves c-Met activation and cellular responsiveness to hepatocyte growth factor. J. Biol. Chem. 279, 26445–26452 https://doi.org/10.1074/jbc.M314254200
- 115 Nakayama, M., Sakai, K., Yamashita, A., Nakamura, T., Suzuki, Y. and Matsumoto, K. (2013) Met/HGF receptor activation is regulated by juxtamembrane Ser985 phosphorylation in hepatocytes. *Cytokine* **62**, 446–452 https://doi.org/10.1016/j.cyto.2013.04.006
- 116 Frampton, G.M., Ali, S.M., Rosenzweig, M., Chmielecki, J., Lu, X., Bauer, T.M. et al. (2015) Activation of MET via diverse exon 14 splicing alterations occurs in multiple tumor types and confers clinical sensitivity to MET inhibitors. *Cancer Discov.* 5, 850–859 https://doi.org/10.1158/2159-8290. CD-15-0285
- 117 Paik, P.K., Drilon, A., Fan, P.D., Yu, H., Rekhtman, N., Ginsberg, M.S. et al. (2015) Response to MET inhibitors in patients with stage IV lung adenocarcinomas harboring MET mutations causing exon 14 skipping. *Cancer Discov.* **5**, 842–849 https://doi.org/10.1158/2159-8290.CD-14-1467
- 118 Kong-Beltran, M., Seshagiri, S., Zha, J., Zhu, W., Bhawe, K., Mendoza, N. et al. (2006) Somatic mutations lead to an oncogenic deletion of met in lung cancer. *Cancer Res.* 66, 283–289 https://doi.org/10.1158/0008-5472.CAN-05-2749
- 119 Abella, J.V., Peschard, P., Naujokas, M.A., Lin, T., Saucier, C., Urbe, S. et al. (2005) Met/Hepatocyte growth factor receptor ubiquitination suppresses transformation and is required for Hrs phosphorylation. *Mol. Cell. Biol.* **25**, 9632–9645 https://doi.org/10.1128/MCB.25.21.9632-9645.2005
- 120 Lu, X., Peled, N., Greer, J., Wu, W., Choi, P., Berger, A.H. et al. (2017) MET exon 14 mutation encodes an actionable therapeutic target in lung adenocarcinoma. *Cancer Res.* 77, 4498–4505 https://doi.org/10.1158/0008-5472.CAN-16-1944
- 121 Vigna, E., Gramaglia, D., Longati, P., Bardelli, A. and Comoglio, P.M. (1999) Loss of the exon encoding the juxtamembrane domain is essential for the oncogenic activation of TPR-MET. *Oncogene* **18**, 4275–4281 https://doi.org/10.1038/sj.onc.1202791
- 122 Mak, H.H.L., Peschard, P., Lin, T., Naujokas, M.A., Zuo, D. and Park, M. (2007) Oncogenic activation of the Met receptor tyrosine kinase fusion protein, Tpr-Met, involves exclusion from the endocytic degradative pathway. *Oncogene* **26**, 7213–7221 https://doi.org/10.1038/sj.onc.1210522
- 123 Bardelli, A., Longati, P., Williams, T.A., Benvenuti, S. and Comoglio, P.M. (1999) A peptide representing the carboxyl-terminal tail of the met receptor inhibits kinase activity and invasive growth. J. Biol. Chem. 274, 29274–29281 https://doi.org/10.1074/jbc.274.41.29274
- 124 Gual, P., Giordano, S., Anguissola, S. and Comoglio, P.M. (2001) Differential requirement of the last C-terminal tail of Met receptor for cell transformation and invasiveness. *Oncogene* **20**, 5493–5502 https://doi.org/10.1038/sj.onc.1204713
- 125 Yokoyama, N., Ischenko, I., Hayman, M.J. and Miller, W.T. (2005) The C terminus of RON tyrosine kinase plays an autoinhibitory role. J. Biol. Chem. 280, 8893–8900 https://doi.org/10.1074/jbc.M412623200
- 126 Sternberg, M.J. and Gullick, W.J. (1990) A sequence motif in the transmembrane region of growth factor receptors with tyrosine kinase activity mediates dimerization. *Protein Eng.* **3**, 245–248 https://doi.org/10.1093/protein/3.4.245
- 127 Teese, M.G. and Langosch, D. (2015) Role of gxxxG motifs in transmembrane domain interactions. *Biochemistry* **54**, 5125–5135 https://doi.org/10. 1021/acs.biochem.5b00495
- 128 Lemmon, M.A., Treutlein, H.R., Adams, P.D., Brunger, A.T. and Engelman, D.M. (1994) A dimerization motif for transmembrane alpha-helices. *Nat. Struct. Biol.* **1**, 157–163 https://doi.org/10.1038/nsb0394-157
- 129 Finger, C., Escher, C. and Schneider, D. (2009) The single transmembrane domains of human receptor tyrosine kinases encode self-interactions. *Sci. Signal.* **2**, ra56 https://doi.org/10.1126/scisignal.2000547